

Study of Semileptonic Decays $B^\pm \rightarrow \eta^{(\prime)} l \nu$

C. S. Kim^{a*} and Ya-Dong Yang^{b†}

^a *Department of Physics and IPAP, Yonsei University, Seoul 120-749, Korea*

^b *Department of Physics, Technion-Israel Institute of Technology, Haifa 32000, Israel*

February 1, 2008

Abstract

We study semileptonic decays $B^\pm \rightarrow \eta^{(\prime)} l \nu$, which are suggested to be used to extract the hadronic form factors of B meson decays to $\eta(\eta')$ and the angle of $\eta - \eta'$ mixing. This would be of great benefit to theoretical studies of B nonleptonic decays involving η and η' , and could lead to a reliable and complementary determination of V_{ub} . The branching ratios are estimated to be $\mathcal{B}(B^\pm \rightarrow \eta^{(\prime)} l \nu) = 4.32 \pm 0.83$ (2.10 ± 0.40) $\times 10^{-5}$, which could be extensively studied experimentally at *BaBar* and Belle.

PACS Numbers: 13.20He, 12.15Hh

* cskim@mail.yonsei.ac.kr

† yang@physics.technion.ac.il

Semileptonic B decays are subjects of considerable interests that have been extensively studied with applications of various nonperturbative theoretical frameworks. They offer the most direct method to determine the weak mixing angles and to probe strong interaction confinement phenomenology of hadronic transitions. Recently V_{cb} has been determined from semileptonic B decays and becomes the third most accurately measured Cabibbo-Kobayashi-Maskawa (CKM) matrix element [1]. CLEO Collaborations [2] have made measurements of the decays $B^0 \rightarrow \pi^- l^+ \nu$ and $\rho^- l^+ \nu$ with the results

$$\begin{aligned}\mathcal{B}(B^0 \rightarrow \pi^- l^+ \nu) &= (1.8 \pm 0.4 \pm 0.3 \pm 0.2) \times 10^{-4}, \\ \mathcal{B}(B^0 \rightarrow \rho^- l^+ \nu) &= (2.57 \pm 0.29_{-0.46}^{+0.33} \pm 0.41) \times 10^{-4} \\ \text{and } |V_{ub}| &= (3.25 \pm 0.14_{-0.29}^{+0.21} \pm 0.55) \times 10^{-3}.\end{aligned}$$

It is known that extracting $|V_{ub}|$ from the measured decay rates require significant inputs from theoretical estimations of the hadronic form factors which involve complex strong-interaction dynamics. With *BaBar* and Belle taking data, we are entering a new era of B physics. Prospects for accurate measurement of these decay modes become excellent. We can foresee that the decays $B^\pm \rightarrow \eta l \nu$ and $\eta' l \nu$ could be also observed at B factories in the near future. In this Brief Report, we study the decays $B^\pm \rightarrow \eta l \nu$ and $\eta' l \nu$ to show that many interesting physical observables can be extracted from measurements of these decays.

Amplitudes of exclusive semileptonic $B \rightarrow Pl \nu$ ($l = \mu, e$ and $P = \pi, \eta, \eta'$) can be written as

$$\mathcal{M}(B \rightarrow Pl \nu) = \frac{G_F}{\sqrt{2}} V_{ub} \bar{l} \gamma_\mu (1 - \gamma_5) \nu \langle P(p_P) | \bar{u} \gamma^\mu (1 - \gamma_5) b | B(p_B) \rangle, \quad (1)$$

where the hadronic transition matrix can be parameterized as

$$\langle P(p_P) | \bar{u} \gamma_\mu (1 - \gamma_5) b | B(p_B) \rangle = F_+^{B \rightarrow P}(q^2) (p_B + p_P)_\mu + F_-^{B \rightarrow P}(q^2) (p_B - p_P)_\mu. \quad (2)$$

Here, $q = p_B - p_P$, and $F_{+(-)}^{B \rightarrow P}(q^2)$ are the relevant form factors. Using these notations, the double differential decay width is

$$\frac{d\Gamma(B \rightarrow Pl \nu)}{dE_l dq^2} = G_F^2 |V_{ub}|^2 \frac{1}{16\pi^3 M_B} \left| F_+^{B \rightarrow P}(q^2) \right|^2 \left[2E_l (m_B^2 + q^2 - m_P^2) - m_B (4E_l^2 + q^2) \right], \quad (3)$$

where we have neglected the lepton mass.

To calculate the semileptonic decay width, we have to know precisely the form factors $F_+^{B \rightarrow P}(q^2)$, which challenge our poor knowledge of nonperturbative QCD. In recent years, considerable progress has been made in the calculations of $F_+^{B \rightarrow \pi}(q^2)$ with various theoretical approaches: quark models [3], QCD sum rules [4, 5] and lattice QCD [6, 7]. Combining the results of different approaches, say, predictions of QCD sum rules in low q^2 region and of lattice QCD in high q^2 region, we could possibly obtain a good theoretical description of $F_+^{B \rightarrow \pi}(q^2)$ in the whole q^2 region. However, both QCD sum rule and lattice calculations of the form factors $F_+^{B \rightarrow \eta^{(\prime)}}(q^2)$ are not yet available in the literature. Therefore, we will use $SU(3)_F$ symmetry to relate them to $F_+^{B \rightarrow \pi}(q^2)$. For $\eta - \eta'$ mixing, we adopt the scheme [8, 9, 10]

$$\begin{aligned} |\eta\rangle &= \cos \phi |\eta_q\rangle - \sin \phi |\eta_s\rangle, \\ |\eta'\rangle &= \sin \phi |\eta_q\rangle + \cos \phi |\eta_s\rangle, \end{aligned} \quad (4)$$

where $|\eta_q\rangle = (u\bar{u} + d\bar{d})/\sqrt{2}$, $|\eta_s\rangle = s\bar{s}$, and $\phi = 39.3^\circ$ is the fitted mixing angle [9]. Assuming $SU(3)_F$ symmetry, the form factors $F_+^{B \rightarrow \eta^{(\prime)}}(q^2)$ are related to $F_+^{B \rightarrow \pi}(q^2)$ by the relations

$$F_+^{B \rightarrow \eta}(q^2) = \cos \phi F_+^{B \rightarrow \pi}(q^2), \quad F_+^{B \rightarrow \eta'}(q^2) = \sin \phi F_+^{B \rightarrow \pi}(q^2). \quad (5)$$

The form factor $F_+^{B \rightarrow \pi}(q^2)$ is known to be dominated by B^* pole in the small-recoil region $q^2 \sim (m_B - m_\pi)^2$ and to scale as $F_+^{B \rightarrow \pi}(q^2 \simeq m_B^2) \sim \sqrt{m_B}$ in the heavy quark limit [11]. Recent studies [7, 12, 13] imply the dipole behavior for $F_+(q^2)$ in the large-recoil region $q^2 \sim 0$. The easiest way to extrapolate the q^2 dependence is to suppose the dipole behavior for $F_+(q^2)$ [6, 14]

$$F_+^{B \rightarrow \pi}(q^2) = \frac{F_+^{B \rightarrow \pi}(0)}{(1 - \frac{q^2}{m_{B^*}^2})^2}, \quad (\text{Dipole}) \quad (6)$$

where m_{B^*} is the pole mass of $B^*(1^-)$ associated with the weak current induced by the decay.

Becirevic and Kaidalov (BK) [15] have also proposed a numerical parameterization which satisfies the heavy quark scaling laws [11] and most of the known constraints [15],

$$F_+^{B \rightarrow \pi}(q^2) = \frac{c_B(1 - \alpha_{B\pi})}{(1 - \frac{q^2}{m_{B^*}^2})(1 - \alpha_{B\pi} \frac{q^2}{m_{B^*}^2})}. \quad (7)$$

We can read from here $F_+^{B \rightarrow \pi}(0) = c_B(1 - \alpha_{B\pi})$. Using BK parameterization to fit their light-cone QCD sum rule (LCSR) calculations, Khodjamirian *et al.* found $\alpha_{B\pi} = 0.32_{-0.07}^{+0.21}$ and

$F_+^{B \rightarrow \pi}(0) = 0.28 \pm 0.05$ [4]. The recent results from Lattice QCD are [16]

$$\alpha_{B\pi} = 0.40 \pm 0.15, \quad F_+^{B \rightarrow \pi}(0) = 0.26 \pm 0.05, \quad (\text{Lattice I})$$

$$\alpha_{B\pi} = 0.45 \pm 0.17, \quad F_+^{B \rightarrow \pi}(0) = 0.28 \pm 0.06, \quad (\text{Lattice II}) \quad (8)$$

where the two sets of results (Lattice I, II) correspond to two different methods used in Ref. [16].

To eliminate the effect of large uncertainty in V_{ub} , we relate the branching ratios $B \rightarrow \eta^{(\prime)} l \nu$ to $\mathcal{B}(B^- \rightarrow \pi^0 l \nu)$, and get

$$\begin{aligned} \mathcal{R}_1 = \frac{\mathcal{B}(B^- \rightarrow \eta l \nu)}{\mathcal{B}(B^- \rightarrow \pi^0 l \nu)} &= |\cos \phi|^2 \frac{\int_0^{(m_B - m_\eta)^2} dq^2 |F_+^{B \rightarrow \pi}(q^2)|^2 \left((m_B^2 + m_\eta^2 - q^2)^2 - 4m_B^2 m_\eta^2 \right)^{\frac{3}{2}}}{\int_0^{(m_B - m_\pi)^2} dq^2 |F_+^{B \rightarrow \pi}(q^2)|^2 \left((m_B^2 + m_\pi^2 - q^2)^2 - 4m_B^2 m_\pi^2 \right)^{\frac{3}{2}}} \\ &= |\cos \phi|^2 \times \begin{cases} 0.527, & (\text{Dipole}) \\ 0.813, & (\text{LCSR}) \\ 0.802, & (\text{Lattice I}) \\ 0.794, & (\text{Lattice II}) \end{cases} \quad (9) \end{aligned}$$

$$\begin{aligned} \mathcal{R}_2 = \frac{\mathcal{B}(B^- \rightarrow \eta' l \nu)}{\mathcal{B}(B^- \rightarrow \pi^0 l \nu)} &= |\sin \phi|^2 \frac{\int_0^{(m_B - m_{\eta'})^2} dq^2 |F_+^{B \rightarrow \pi}(q^2)|^2 \left((m_B^2 + m_{\eta'}^2 - q^2)^2 - 4m_B^2 m_{\eta'}^2 \right)^{\frac{3}{2}}}{\int_0^{(m_B - m_\pi)^2} dq^2 |F_+^{B \rightarrow \pi}(q^2)|^2 \left((m_B^2 + m_\pi^2 - q^2)^2 - 4m_B^2 m_\pi^2 \right)^{\frac{3}{2}}} \\ &= |\sin \phi|^2 \times \begin{cases} 0.310, & (\text{Dipole}) \\ 0.599, & (\text{LCSR}) \\ 0.584, & (\text{Lattice I}) \\ 0.573, & (\text{Lattice II}). \end{cases} \quad (10) \end{aligned}$$

Using the CLEO result [2], $\mathcal{B}(B^0 \rightarrow \pi^+ l \nu) = (1.8 \pm 0.6) \times 10^{-4}$, and the relations $\mathcal{B}(B^0 \rightarrow \pi^+ l \nu) = 2\mathcal{B}(B^- \rightarrow \pi^0 l \nu)$, $\phi = 39.3^\circ$, we get

$$\mathcal{B}(B^- \rightarrow \eta l \nu) = \begin{cases} (2.84 \pm 0.95) \times 10^{-5}, & (\text{Dipole}) \\ (4.38 \pm 1.46) \times 10^{-5}, & (\text{LCSR}) \\ (4.32 \pm 1.44) \times 10^{-5}, & (\text{Lattice I}) \\ (4.28 \pm 1.42) \times 10^{-5}, & (\text{Lattice II}), \end{cases} \quad (11)$$

$$\mathcal{B}(B^- \rightarrow \eta' l \nu) = \begin{cases} (1.12 \pm 0.37) \times 10^{-5}, & (\text{Dipole}) \\ (2.16 \pm 0.72) \times 10^{-5}, & (\text{LCSR}) \\ (2.10 \pm 0.70) \times 10^{-5}, & (\text{Lattice I}) \\ (2.06 \pm 0.68) \times 10^{-5}, & (\text{Lattice II}). \end{cases} \quad (12)$$

We can see that the predictions of LCSR form factors [4] agree very well with those of Lattice (I, II) QCD [16]. Averging predictions from Lattice QCD and LCSR, we obtain

$$\begin{aligned}\mathcal{B}(B^- \rightarrow \eta l \nu) &= (4.32 \pm 0.83) \times 10^{-5}, \\ \mathcal{B}(B^- \rightarrow \eta' l \nu) &= (2.10 \pm 0.40) \times 10^{-5}.\end{aligned}\tag{13}$$

We note that the ratios $\mathcal{R}_1$ and $\mathcal{R}_2$ are independent of the value of $F_+^{B \rightarrow \pi}(0)$, but very sensitive to the details of its q^2 dependence. We also note that to give the same numerical predictions for $B \rightarrow \pi l \nu$, the $F_+^{B \rightarrow \pi}(0)$ for dipole parameterization should be smaller than that for BK parameterization. If the same value $F_+^{B \rightarrow \pi}(0)$ is used in both BK and dipole parameterizations, one will find

$$\mathcal{R}_3 = \frac{\mathcal{B}^{\text{dipole}}(B^- \rightarrow \pi^0 l \nu)}{\mathcal{B}^{\text{LCSR}}(B^- \rightarrow \pi^0 l \nu)} = 3.13,\tag{14}$$

which implies that the dipole form factor will overestimate the decay rates because π meson is very light and the lepton pair invariant mass q^2 can be very near the B_u^* pole. Therefore, theoretical predictions for $B \rightarrow \pi l \nu$ (and $B \rightarrow \eta^{(\prime)} l \nu$ in turn) are very sensitive to the q^2 dependence of $F_+^{B \rightarrow \pi}(q^2)$. It is well known that the extraction of V_{ub} from decay rates of $B \rightarrow \pi(\rho) l \nu$ suffers from large theoretical uncertainties in the hadronic form factors. Testing the predictions and eventual measurements of $d\Gamma/dq^2$ can provide valuable information on the hadronic form factors governing $b \rightarrow u l \nu$ decays, and hence lead to a reliable determination of V_{ub} . With much more data to arrive soon from B factories, the q^2 and the lepton energy distributions can be precisely measured and be used to distinguish these form factor parameterizations, and to extract V_{ub} . The determination of V_{ub} from $B \rightarrow \eta^{(\prime)} l \nu$ would represent a powerful method complementary to the determination of V_{ub} from other exclusive decay modes, e.g., from $B \rightarrow \pi(\rho) l \nu$.

In Fig. 1, we plot the q^2 distributions and the lepton energy E_l distributions of the decays $B^- \rightarrow \eta^{(\prime)} l \nu$, where we have normlized the form factors to give $\mathcal{B}(B^- \rightarrow \pi^0 l \nu) = 9 \times 10^{-5}$. We find that both LCSR and Lattice QCD predict very consistent lepton energy distributions as well as consistent decay rates for the decays. Integrating out the lepton energy in Eq. (3), one obtains

$$\frac{d\Gamma(B \rightarrow P l \nu)}{dq^2} = \frac{G_F^2 |V_{ub}|^2}{192\pi^3 M_B^3} \left| F_+^{B \rightarrow P}(q^2) \right|^2 \left[(m_B^2 + m_P^2 - q^2)^2 - 4m_B^2 m_P^2 \right]^{\frac{3}{2}}.\tag{15}$$

At maximum recoil point ($q^2 = 0$), we have

$$\mathcal{R}_4 = \frac{d\Gamma(B \rightarrow \eta^{(\prime)} l \nu)/dq^2}{d\Gamma(B^- \rightarrow \pi^0 l \nu)/dq^2} \Big|_{q^2=0} = \frac{(m_B^2 - m_{\eta^{(\prime)}}^2)^3}{(m_B^2 - m_\pi^2)^3} \left| \frac{F_+^{B \rightarrow \eta^{(\prime)}}(0)}{F_+^{B \rightarrow \pi^0}(0)} \right|^2, \quad (16)$$

$$\begin{aligned} \mathcal{R}_5 &= \frac{d\Gamma(B \rightarrow \eta' l \nu)/dq^2}{d\Gamma(B^- \rightarrow \eta l \nu)/dq^2} \Big|_{q^2=0} = \frac{(m_B^2 - m_{\eta'}^2)^3}{(m_B^2 - m_\eta^2)^3} \left| \frac{F_+^{B \rightarrow \eta^{(\prime)}}(0)}{F_+^{B \rightarrow \eta}(0)} \right|^2 \\ &= \frac{(m_B^2 - m_{\eta'}^2)^3}{(m_B^2 - m_\eta^2)^3} |\cot \phi|^2. \end{aligned} \quad (17)$$

As indicated by QCD sum rule calculations [4, 5], the value of $F_+^{B \rightarrow \eta^{(\prime)}}(q^2)$ is rather stable under the variation of q^2 when the value of q^2 is small. So the ratios $\mathcal{R}_4$ and $\mathcal{R}_5$ can be safely extrapolated to few GeV^2 to make the phase spaces sizable. Once the ratios are measured, they can be used to extract the form factor $F_+^{B \rightarrow \eta^{(\prime)}}(0)$ and the mixing angle ϕ from the above relations.

In the literature, semileptonic decays $D_s \rightarrow \eta(\eta') l \nu$ have been taken as sources of extracting η – η' mixing angle and testing the mixing schemes [9, 17]. We note that the decays $D_s \rightarrow \eta(\eta') l \nu$ involve strange contents $|\eta_s\rangle$ of $\eta(\eta')$, and $B^- \rightarrow \eta(\eta') l \nu$ involve non-strange contents $|\eta_q\rangle$ of $\eta(\eta')$, therefore $B^- \rightarrow \eta(\eta') l \nu$ and $D_s \rightarrow \eta(\eta') l \nu$ could provide combined testing of η – η' mixing scheme.

As it is well known that η and η' are too complicated objects to be reliably described within QCD yet, it may be very hard to calculate the transition form factors $F_+^{B \rightarrow \eta^{(\prime)}}(q^2)$ within the frameworks of lattice QCD and QCD sum rules. The experimental extraction of those form factors will improve our theoretical understanding of many interesting nonleptonic B decay modes involving η and η' , and might shed light on the problem, currently under discussion [18], of the puzzling large branching ratios of $B \rightarrow K \eta'$ observed by CLEO [19].

Finally, we note a few experimental comments: Background for $B \rightarrow \eta^{(\prime)} l \nu$ would be much smaller than that of $B \rightarrow \pi l \nu$, due to much lower multiplicity, since the random background caused by $B \rightarrow \eta X$ is about an order of magnitude smaller than that by $B \rightarrow \pi X$. The reconstruction of $\eta \rightarrow \gamma\gamma$ in experimental analyses may be much easier than $\pi^0 \rightarrow \gamma\gamma$, even though the signal/noise ratio is worse, because the mass of η is much bigger than that of π^0 . And we could even require the momentum of η to be bigger than 1 GeV to remove combinatorial backgrounds substantially.

To conclude, we studied semileptonic decays $B^\pm \rightarrow \eta^{(\prime)} l \nu$, which can be used to extract the

hadronic form factors of B meson decays to $\eta(\eta')$ and the angle of $\eta - \eta'$ mixing. The branching ratios are found to be $\mathcal{B}(B^- \rightarrow \eta^{(\prime)} l \nu) = 4.32 \pm 0.83 \text{ } (2.10 \pm 0.40) \times 10^{-5}$.

Acknowledgments

We thank G. Cvetic, Hongjoo Kim and Y.J. Kwon for careful reading of the manuscript and their valuable comments. The work of C.S.K. was supported by Grant No. 2000-015-DP0077 of the KRF. Y.Y is supported by the US-Israel Binational Science Foundation and the Israel Science Foundation.

References

- [1] Partical Data Group, C. Caso *et al.*, Eur. Phys. J. **C15**, 1 (2000).
- [2] CELO Collaboration: J.P. Alexander *et al.*, Phys. Rev. Lett. **77**, 5000 (1996); Phys. Rev. **D61**, 052001 1(2000).
- [3] M. Wirbel, B. Stech and M. Bauer, Z. Phys. **C29**, 637 (1985); N. Isgur, D. Scora, B. Grinstein and M.B Wise, Phys. Rev.**D39**, 799 (1989); D. Scora and N. Isgur, Phys. Rev. **D52**, 2783 (1995); I. L. Grach, I. M. Narodetskii, and S. Simula, Phys. Lett, **B385**, 317 (1996); M. Bayer and D. Melikhov, Phys. Lett. **B436**, 344 (1998); P. Colangelo and P.Santorelli, Phys. Lett. **B327**, 123(1994); M. Ladisa, G. Nardulli and P. Santorelli, Phys. Lett, **B455**, 283 (1999); A. Deandrea, R. Gatto, G. Nardulli and A. D. Polosa, Phys. Rev. **D61**, 017502 (2000).
- [4] A. Khodjamirian, R. Rückl, S. Weinzierl, C. Winhart and O. Yakovlev, Phys. Rev. **D62**, 114002 (2000).
- [5] E. Bagan, P. Ball and V. M. Braun, Phys. Lett. **B417**, 154 (1998); P. Ball, Phys. Rev. **D48**, 3190 (1993); JHEP **09**, 005 (1998).
- [6] UKQCD Collaboration: K.C. Bowler *et al.*, Phys. Lett. **B486**, 111 (2000).

- [7] A. Abada, *et al.*, Nucl. Phys. **B416**, 675 (1994); C.R. Allton, *et al.*, Phys. Lett. **B345**, 513 (1995).
- [8] J.L. Rosner, Phys. Rev. **D27**, 1101(1983).
- [9] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. **D58**, 114006 (1998); Phys. Lett. B449, 339 (1999); T. Feldmann, Int. J. Mod. Phys. **A15**, 159 (2000) and references therein.
- [10] A. Bramon, R.Escribano and M.D. Scadron, Eur. Phys. J. **C7**, 271(1998).
- [11] N. Isgur and M.B. Wise, Phys. Rev. **D41**, 151 (1990); Phys. Rev. **D42**, 2388 (1990).
- [12] J. Charles, A. Le Yaouanc, L. Oliver, O. Pene and J. C. Raynal, Phys. Rev. **D60**, 014001 (1999).
- [13] H. Y.Cheng, C. Y. Cheung and C. W. Hwang, Phys. Rev. **D55**, 1559 (1997).
- [14] R. Alexan *et al.*, Phys. Rev. **D51**, 6235 (1995); UKQCD Collaborations, D.R. Burford *et al.*, Nucl. Phys. **B447**, 425 (1995)
- [15] D. Becirevic and A. Kaidalov, Phys. Lett. **B478**, 417 (2000).
- [16] A. Abada, D. Becirevic, Ph. Boucaud, J.P. Leroy, V. Lubicz and F. Mescia, hep-ph/0011065.
- [17] D. Melikhov, Phys. Lett. **B394**, 385 (1997).
- [18] Dong-sheng Du, C.S. Kim and Ya-dong Yang, Phys. Lett. **B426**, 133 (1998); A. Ali, J. Chay, C. Greub and P. Ko, Phys. Lett. **B424**, 161 (1998); A. Datta, S. Pakvasa and X.G. He, Phys. Lett. **B419**, 369 (1998); A.S. Dighe, M. Gronau and J.L. Rosner, Phys. Rev. Lett. **79**, 4333 (1997); Maozhi Yang and Ya-dong Yang, hep-ph/0012208, Nucl. Phys. **B609**, 469 (2001).
- [19] CLEO Collaboration: B. H. Behrens *et al.*, Phys. Rev. Lett. **80**, 3710 (1998); S. J. Richichi, *et al.*, Phys. Rev. Lett. **85**, 520 (2000).




Figure 1: The spectra $d\mathcal{B}/dq^2$ as function of q^2 and the spectra $d\mathcal{B}/dE_l$ as function of the electron energy E_l . The thick solid, long-dashed and short-dashed curves are the distributions of $d\mathcal{B}(B \rightarrow \eta l \nu)$ with LCSR [4] and Lattice (I, II) QCD [16] form factors, and the thin curves are those for $d\mathcal{B}(B \rightarrow \eta' l \nu)$.